

# The Friendship Paradox

Barum Park

Department of Sociology  
Cornell University  
b.park@cornell.edu

April 27, 2023

## The Paradox

The [friendship paradox](#) states that, on average, your friends will have more friends than you have. It is called a paradox because most people believe that they have more friends than their friends (is this true?). Interestingly enough, Feld called this the "class size paradox," but his own terminology was not adopted in the literature. Formally, we might state the paradox as follows:

Let  $G(V, E)$  be a connected graph with  $|V| = n$  nodes,  $|E| = m$  edges, and a degree sequence  $\text{Deg}(G) = \{d_1, d_2, \dots, d_n\}$ , where  $d_i$  is the degree of node  $i \in V$ . Let  $\bar{d}$  be the nodes' average number of friends and  $\tilde{d}$  the average number of nodes' friends of friends. Then,  $\tilde{d} \geq \bar{d}$ .

The friendship paradox is, in fact, quite easy to prove. The easiest proof is a "story proof." Suppose all nodes report how many friends their friends have. How many reports are made? Since each edge in the network connects two nodes, and each edge entails two reports, the total number of reports is  $2m$ . Next, consider a focal node  $v$  with degree  $d_v$ . All of  $v$ 's friends will report that  $v$  has  $d_v$  friends. As there are  $d_v$  friends who make the report, there will give a total of  $d_v^2$  reported friends of friends. Summing this over all nodes, we see that  $\sum_{i=1}^n d_i^2$  friends of friends are reported. It follows that the average number of reported friends of friends is

$$\tilde{d} = \frac{1}{2m} \sum_{i=1}^n d_i^2.$$

It remains to show that  $\tilde{d} \geq \bar{d}$ . By the [Cauchy-Schwarz Inequality](#),<sup>1</sup> we have

$$2m\tilde{d} = \sum_{i=1}^n d_i^2 \geq \frac{1}{n} \left( \sum_{i=1}^n d_i \right)^2 = n\bar{d}^2$$

Now, note that  $n\bar{d} = 2m$  (when we sum all degrees, we get two times the total number of edges, since every edge is counted twice). Hence,  $n\bar{d}^2 = 2m\bar{d}$ , and dividing both sides of the inequality by  $2m > 0$  gives

$$\tilde{d} \geq \bar{d}$$

---

<sup>1</sup>The Cauchy-Schwarz Inequality states that for two vectors  $v, w \in \mathbb{R}^n$ ,  $(v \cdot v)(w \cdot w) \geq |v \cdot w|^2$ , where  $\cdot$  denotes the inner (dot) product. Choosing  $v$  to be a vector of 1s and  $w$  to be the degree sequence of  $G$  gives  $n \sum_i d_i^2 \geq (\sum_i d_i)^2$ . The inequality in the text is obtained, then, by dividing both sides by  $n$ .

which completes the proof. Notice that the Cauchy-Schwarz Inequality will hold as an equality if and only if  $G$  is a regular graph, i.e., a graph in which each node has exactly the same degree.<sup>2</sup>

## Average Mean Degree of Friends

Feld goes, in fact, one step further to consider the average mean degree of friends—i.e., the mean degree of friends that each individual observes, averaged over all individuals—and argues that this number will be greater than the mean degree.

To prove this claim, consider the adjacency matrix  $A = (a_{ij})$  of the graph from above. The total number of node  $i$ 's friends' friends is  $t_i = \sum_{j=1}^n a_{ij}d_j$ . As  $i$  has  $d_i = \sum_{j=1}^n a_{ij}$  friends, the average number of  $v_i$ 's friends' of friends is thus

$$\tilde{f}_i = \frac{t_i}{d_i} = \frac{1}{d_i} \sum_{j=1}^n a_{ij}d_j.$$

Notice that  $d_i \geq 1$  for all  $i = 1, 2, \dots, n$  since the graph is assumed to be connected. Hence,  $\tilde{f}_i$  is well-defined. The average of  $\tilde{f}_i$  is

$$\begin{aligned} \bar{f} &= \frac{1}{n} \sum_{i=1}^n \tilde{f}_i = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n a_{ij} \left( \frac{d_j}{d_i} \right) \\ &= \frac{1}{n} \sum_{i,j:(i,j) \in E} \left( \frac{d_j}{d_i} \right) \end{aligned} \tag{1}$$

since  $a_{ij} = 1$  if the edge  $(i, j) \in E$ . Notice that this sum is over all edges  $(i, j) \in E$ , where the contribution of each edge  $(i, j)$  to the total sum is

$$\frac{d_i}{d_j} + \frac{d_j}{d_i}.$$

Compare  $\bar{f}$  with the average degree:

$$\bar{d} = \frac{1}{n} \sum_{i=1}^n d_i = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n a_{ij} = \frac{1}{n} \sum_{i,j:(i,j) \in E} 1, \tag{2}$$

which sums over all edges of  $G$  as well, but where the contribution of each edge  $(i, j)$  is  $1 + 1$ . Hence,  $\bar{d} \leq \bar{f}$  if

$$\frac{d_i}{d_j} + \frac{d_j}{d_i} \geq 2$$

for all  $i, j \in V$ . And, indeed,

$$\frac{d_i}{d_j} + \frac{d_j}{d_i} = 2 + \frac{(d_i - d_j)^2}{d_i d_j} \geq 2$$

since  $d_i > 0$ . So we are done.

As in the original friendship paradox, notice that equation (1) will reduce to (2), when  $d_i = c$  for all  $i$ —i.e., if  $G$  is a regular graph.

## Generalized Friendship Paradox

There is also something called the [generalized friendship paradox](#). This term refers to nothing more than the observation that for any node-level characteristic that is positively correlated with degree, a similar

---

<sup>2</sup>In other words,  $|v \cdot w|^2 = (v \cdot v)(w \cdot w)$  if and only if  $v = \alpha w$ , where  $\alpha$  is a constant. Thus,  $\bar{d} = \tilde{d}$  if and only if the degree sequence and a vector of ones are [linearly dependent](#).

result holds: namely, your friends will have more of that characteristic, on average, than you have. To see why this is true, consider a characteristic  $x$ . The average  $x$  for the nodes and the node's friends is

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad \text{and} \quad \tilde{x} = \frac{\sum_{i=1}^n d_i x_i}{\sum_{i=1}^n d_i}.$$

Notice that if  $x_i = d_i$ , we get back to the original friendship paradox.

We might say that the generalized friendship holds if  $\tilde{x} > \bar{x}$  or, equivalently, when  $\tilde{x} - \bar{x} > 0$ . A bit of algebra reveals that

$$\begin{aligned} \tilde{x} - \bar{x} &= \frac{\sum_{i=1}^n d_i x_i}{\sum_{i=1}^n d_i} - \bar{x} \\ &= \frac{\sum_{i=1}^n (d_i - \bar{d})(x_i - \bar{x}) + n\bar{d}\bar{x}}{n\bar{d}} - \bar{x} \\ &= \frac{\sum_{i=1}^n (d_i - \bar{d})(x_i - \bar{x})}{n\bar{d}} \\ &= \frac{\text{Cov}(d, x)}{\bar{d}}. \end{aligned}$$

As  $\bar{d} > 0$ , this shows that  $\tilde{x} - \bar{x}$  if  $\text{Cov}(x, d) > 0$ . In other words, the generalized friendship paradox will hold for any characteristic that is positively correlated with degree.